



P O P U L A T I O N R E F E R E N C E B U R E A U

Female Genital Mutilation/ Cutting: Data and Trends

► ► ► **UPDATE 2010**

An estimated 100 million to 140 million girls and women worldwide have undergone female genital mutilation/cutting (FGM/C) and more than 3 million girls are at risk for cutting each year on the African continent alone.

FGM/C is generally performed on girls between ages 4 and 12, although it is practiced in some cultures as early as a few days after birth or as late as just prior to marriage. Typically, traditional excisors have carried out the procedure, but recently a discouraging trend has emerged in some countries where medical professionals are increasingly performing the procedure.

FGM/C poses serious physical and mental health risks for women and young girls, especially for women who have undergone extreme forms of the procedure (see Box 2 for types of cutting). According to a 2006 WHO study, FGM/C can be linked to increased complications in childbirth and even maternal deaths. Other side effects include severe pain, hemorrhage, tetanus, infection, infertility, cysts and abscesses, urinary incontinence, and psychological and sexual problems.

FGM/C is practiced in at least 28 countries in Africa and a few others in Asia and the Middle East. The 27 developing countries included on this chart are the only ones where data have been systematically collected at this time. FGM/C is practiced at all educational levels and in all social classes and occurs among many religious groups (Muslims, Christians, and animists), although no religion mandates it. Prevalence rates vary significantly from country to country (from nearly 98 percent in Somalia to less than 1 percent in Uganda) and even within countries.

Since the early 1990s, FGM/C has gained recognition as a health and human rights issue among African governments, the international community, women's organizations, and professional associations. Global and national efforts to end FGM/C have supported legislation targeting excisors, medical professionals, and families who perpetuate the practice, but political will and implementation remain an issue.

Some of the data that have been collected in recent years give hope to those working toward the abandonment of FGM/C as they reflect lower levels of cutting among girls ages 15 to 19.

Box 1

Deciphering the Terms: Mutilation, Cutting, or Circumcision?

Female genital mutilation/cutting (FGM/C), also known as female circumcision (FC), female genital cutting (FGC), and female genital mutilation (FGM), involves the cutting or alteration of the female genitalia for social rather than medical reasons.

The term FC was widely used for many years to describe the practice; however, it has been largely abandoned as it implies an analogy with male circumcision. Various communities still use the term FC because it is a literal translation from their own languages. Female genital mutilation/cutting is a far more damaging and invasive procedure than male circumcision. FGM/C is often perceived as a way to curtail premarital sex and preserve virginity.

FGM is the term most commonly used by women's rights and health advocates who wish to emphasize the damage caused by the procedure. In the mid-1990s, many organizations decided to shift to the use of the more neutral term, female genital cutting, because they considered FGM to be judgmental, pejorative, and not conducive to discussion and collaboration on abandonment. The U.S. Agency for International Development currently uses the term FGM/C.

Sources: Anika Rahman and Nahid Toubia, *Female Genital Mutilation: A Guide to Laws and Policies Worldwide* (London and New York: Zed Books, 2000): 4; and *Abandoning Female Genital Mutilation/Cutting: Information From Around the World* (Washington, DC: Population Reference Bureau, 2005).

Box 2

Types of Female Genital Mutilation/Cutting

Female genital mutilation/cutting (FGM/C) refers to a variety of operations involving partial or total removal of female external genitalia. The female external genital organ consists of the vulva, which is comprised of the labia majora, labia minora, and the clitoris covered by its hood in front of the urinary and vaginal openings.

In 2007, the World Health Organization classified FGM/C into four broad categories:

Type 1 or Clitoridectomy: Partial or total removal of the clitoris and/or the clitoral hood.

Type 2 or Excision: Partial or total removal of the clitoris and the labia minora, with or without excision of the labia majora.

Type 3 or Infibulation: Narrowing of the vaginal orifice with creation of a covering

seal by cutting and placing together the labia minora and/or the labia majora, with or without excision of the clitoris.


Type 4 or Unclassified: All other harmful procedures to the female genitalia for nonmedical purposes, for example, pricking, piercing, incising, scraping, and cauterization.

Note: Current questionnaires used in the Demographic and Health Surveys do not differentiate between Types I and II, but only between whether a girl or woman has been cut, whether tissue has been removed, and whether tissue has been sewn closed.

Source: World Health Organization, *Eliminating Female Genital Mutilation: An Interagency Statement*, OHCHR, UNAIDS, UNDP, UNECA, UNESCO, UNFPA, UNHCR, UNICEF, UNIFEM, WHO (Geneva: WHO, 2008): 23.

Prevalence of FGM/C Among Younger and Older Women


While in some countries there is little difference in prevalence between older women (ages 35 to 39) and younger women (ages 15 to 19), in others—such as Ethiopia, Côte d'Ivoire, and Kenya—the difference is significant. This may be a sign that the practice is being abandoned.



Trends in FGM/C Prevalence


Over the last decade, a downward trend in percent of women cut in some countries indicates that abandonment of FGM/C seems to be taking hold, although in others there still is little or no apparent change.

Percent of women ages 15–49, by survey year



Variations Within and Across Borders

Looking only at national prevalence rates can hide the regional variations within a country. FGM/C often reflects ethnicity or social interactions of communities across national borders.



Female Genital Mutilation/Cutting: Data and Trends

	Data Source		Prevalence by Age (%)		
			15-49	15-19	35-39
Benin	DHS	2006	12.9	7.9	16.3
Burkina Faso	MICS	2006	72.5	59.7	79.8
Cameroon	DHS	2004	1.4	0.4	1.2
Central African Rep.	MICS	2008	25.7	18.7	29.8
Chad	DHS	2004	44.9	43.4	46.2
Côte d'Ivoire	MICS	2006	36.4	28.0	43.8
Djibouti	MICS	2006	93.1	—	—
Egypt	DHS	2008	91.1	80.7	96.4
Eritrea	DHS	2002	88.7	78.3	92.6
Ethiopia	DHS	2005	74.3	62.1	81.2
Gambia	MICS	2005/06	78.3	79.9	79.5
Ghana	MICS	2006	3.8	1.4	5.7
Guinea	DHS	2005	95.6	89.3	98.6
Guinea-Bissau	MICS	2006	44.5	43.5	48.6
Kenya	DHS	2008-09 ^c	27.1	14.6	35.1
Liberia	DHS	2007	58.2	35.9	66.7
Mali	DHS	2006	85.2	84.7	84.9
Mauritania	MICS	2007	72.2	68.0	75.4
Niger	DHS	2006	2.2	1.9	2.9
Nigeria	DHS	2008 ^d	29.6	21.7	33.9
Senegal	DHS	2005	28.2	24.8	30.5
Sierra Leone	MICS	2006	94.0	81.1	97.5
Somalia	MICS	2006	97.9	96.7	98.9
Tanzania	DHS	2004/05	14.6	9.1	16.0
Togo	MICS	2006	5.8	1.3	9.4
Uganda	DHS	2006	0.6	0.5	0.8
Yemen	PAPFAM	2003	38.2	—	—

Female Genital Mutilation/Cutting: Data and Trends

	Data Source		Prevalence by Geographic Area (%)			
			Urban	Rural	Lowest Region	Highest Region
Benin	DHS	2006	9.3	15.4	0.1	58.8
Burkina Faso	MICS	2006	76.0	71.2	—	—
Cameroon	DHS	2004	0.9	2.1	0.0	5.4
Central African Rep.	MICS	2008	20.9	29.3	5.5	76.9
Chad	DHS	2004	47.0	44.4	3.5	92.2
Côte d'Ivoire	MICS	2006	33.9	38.9	12.6	88.0
Djibouti	MICS	2006	93.1	95.5	—	—
Egypt	DHS	2008	85.1	95.5	66.3	92.9
Eritrea	DHS	2002	86.4	90.5	81.5	97.7
Ethiopia	DHS	2005	68.5	75.5	27.1	97.3
Gambia	MICS	2005/06	72.2	82.8	44.8	99.0
Ghana	MICS	2006	1.7	5.7	0.5	56.1
Guinea	DHS	2005	93.9	96.4	86.4	99.8
Guinea-Bissau	MICS	2006	39.0	48.2	28.7	92.7
Kenya	DHS	2008-09 ^c	16.5	30.6	0.8	97.5
Liberia	DHS	2007	39.5	72.0	0.9	86.6
Mali	DHS	2006	80.9	87.4	0.9	98.3
Mauritania	MICS	2007	59.7	84.1	26.6	98.1
Niger	DHS	2006	2.1	2.3	0.1	12.0
Nigeria	DHS	2008 ^d	36.8	25.6	2.7	53.4
Senegal	DHS	2005	21.7	34.4	1.8	93.8
Sierra Leone	MICS	2006	86.4	97.0	80.8	97.0
Somalia	MICS	2006	97.1	98.4	94.4	99.2
Tanzania	DHS	2004/05	7.2	17.6	0.8	57.6
Togo	MICS	2006	4.1	7.3	1.0	22.7
Uganda	DHS	2006	0.2	0.7	0.1	2.4
Yemen	PAPFAM	2003	33.1	40.7	—	—

Female Genital Mutilation/Cutting: Data and Trends

	Data Source		Types of FGM/C (%)		
			Nicked, No Flesh Removed	Flesh Removed	Sewn Closed
Benin	DHS	2006	0.6 ^a	97.5 ^a	3.9 ^{a, b}
Burkina Faso	MICS	2006	0.7	89.5	3.8
Cameroon	DHS	2004	4.0	85.0	5.0
Central African Rep.	MICS	2008	—	92.4	3.3
Chad	DHS	2004	19.4	74.7	2.4
Côte d'Ivoire	MICS	2006	6.1	80.0	5.7
Djibouti	MICS	2006	24.9	6.4	67.2
Egypt	DHS	2008	—	—	—
Eritrea	DHS	2002	46.0	4.1	38.6
Ethiopia	DHS	2005	—	—	6.1
Gambia	MICS	2005/06	—	—	—
Ghana	MICS	2006	—	—	—
Guinea	DHS	2005	1.7	86.4	9.3
Guinea-Bissau	MICS	2006	0.8	91.7	3.2
Kenya	DHS	2008-09 ^c	—	—	—
Liberia	DHS	2007	—	—	—
Mali	DHS	2006	3.0	75.8	10.2
Mauritania	MICS	2007	—	73.9	3.2
Niger	DHS	2006	0.8 ^{a, b}	90.5 ^{a, b}	13.3 ^{a, b}
Nigeria	DHS	2008 ^d	3.0	45.4	5.3
Senegal	DHS	2005	0.2	82.7	11.9
Sierra Leone	MICS	2006	—	—	—
Somalia	MICS	2006	1.3	15.2	79.3
Tanzania	DHS	2004/05	1.9	91.3	2.0
Togo	MICS	2006	7.7	85.7	1.7
Uganda	DHS	2006	—	—	—
Yemen	PAPFAM	2003	—	—	—

Female Genital Mutilation/Cutting: Data and Trends

	Data Source		Traditionally Performed (%)	Medically Performed (%)	National Law
Benin	DHS	2006	99.0	0.6	●
Burkina Faso	MICS	2006	—	—	●
Cameroon	DHS	2004	89.0	4.0	◐
Central African Rep.	MICS	2008	—	—	●
Chad	DHS	2004	94.2	2.7	◐
Côte d'Ivoire	MICS	2006	95.2 ^b	0.5 ^b	●
Djibouti	MICS	2006	—	—	●
Egypt	DHS	2008	66.3	31.9	●
Eritrea	DHS	2002	94.5	0.6	●
Ethiopia	DHS	2005	—	—	●
Gambia	MICS	2005/06	—	—	○
Ghana	MICS	2006	—	—	●
Guinea	DHS	2005	88.7	10.0	●
Guinea-Bissau	MICS	2006	—	—	◐
Kenya	DHS	2008-09 ^c	—	—	●
Liberia	DHS	2007	—	—	○
Mali	DHS	2006	91.7	2.5	◐
Mauritania	MICS	2007	—	—	●
Niger	DHS	2006	97.0	0.5	●
Nigeria	DHS	2008 ^d	73.6	8.9	○
Senegal	DHS	2005	92.5	0.6	●
Sierra Leone	MICS	2006	—	—	○
Somalia	MICS	2006	—	—	○
Tanzania	DHS	2004/05	89.1 ^b	2.0 ^b	●
Togo	MICS	2006	—	—	●
Uganda	DHS	2006	—	—	●
Yemen	PAPFAM	2003	—	—	○

Definitions and Notes

Medically Performed refers to FGM/C performed by a health professional including doctors, nurses, and midwives.

Traditionally Performed refers to FGM/C performed by a traditional practitioner including local specialists known for performing circumcisions, traditional birth attendants, and older women without further designation.

National Laws: ● = Laws that specifically prohibit the practice of FGM/C; ○ = No laws; ◐ = No specific laws, but existing general provisions of criminal codes have been or can be applied to FGM/C.

— Data not available.

^a Total for types of FGM/C adds to more than 100 percent due to multiple responses.

^b Special tabulations by PRB staff.

^c Preliminary report.

^d FGM/C data from the 2008 Nigeria Demographic and Health Survey should be used with caution as the definition of FGM/C was not used consistently by interviewers across regions.

Sources

Main survey sources: Demographic and Health Surveys (DHS); Multiple Indicator Cluster Surveys (MICS), UNICEF; Pan-Arab Project for Family Health (PAPFAM).

Other sources:

Center for Reproductive Rights, *Female Genital Mutilation (FGM): Legal Prohibitions Worldwide* (New York, NY: 2008).

Inter-African Committee on Traditional Practices, *International, Regional and Legal Instruments for the Protection of Women and Girls Against FGM*, accessed at www.iac-ciaf.net on Jan. 6, 2010.

Inter-Parliamentary Union, *Legislation and Other National Provisions*, accessed at www.ipu.org/wmn-e/fgm-prov-b.htm, on Jan. 8, 2010.

Population Reference Bureau, *Abandoning Female Genital Mutilation/Cutting: Information From Around the World*, a CD-ROM (Washington, DC: PRB, 2005).

UNICEF, *Female Genital Mutilation/Cutting: A Statistical Exploration* (November 2005).

Women's United Nations Report Network, *FGM Legislation for 25 African Countries—Female Genital Mutilation*, accessed at http://wunrn.com/news/2006/11_06_06/111206_fgm_legislation.htm, on Jan. 8, 2010.

World Health Organization, *Eliminating Female Genital Mutilation: An Interagency Statement* (Geneva: WHO, 2008).

World Health Organization, *Female Genital Mutilation, Fact Sheet No. 241* (Geneva: WHO, May 2008).

P. Stanley Yoder and Shane Khan, *Numbers of Women Circumcised in Africa: The Production of a Total* (Calverton, MD: ORC Macro, March 2008).

P. Stanley Yoder, N. Abderrahim, and A. Zhuzhuni, *Female Genital Cutting in the Demographic and Health Surveys: A Critical and Comparative Analysis, DHS Comparative Reports No. 7* (Calverton, MD: ORC Macro, September 2004).

Acknowledgments

Authors: Charlotte Feldman-Jacobs and Donna Clifton

Special thanks to Ellen Carnevale, Jay Gribble, Sarah Harbison, Sandra Jordan, Toshiko Kaneda, Kelvin Pollard, Richard Skolnik, and Rhonda Smith.

Cover photo: Pep Bonet/Noor Images

Funding for this publication was provided by the U.S. Agency for International Development, under the BRIDGE Project (Cooperative Agreement GPO-A-00-03-00004-00). The contents are the responsibility of the Population Reference Bureau and do not necessarily reflect the views of USAID or the United States Government.



USAID
FROM THE AMERICAN PEOPLE

BRIDGE



To order copies, contact PRB, 1875 Connecticut Ave., NW, Suite 520, Washington, DC 20009-5728; 202-483-1100; Fax: 202-328-3937; E-mail: prborders@prb.org; www.prb.org.

© 2010 Population Reference Bureau. All rights reserved.